GRUPPO NAZIONALE PER LA DIFESA DALLE CATASTROFI IDROGEOLOGICHE – U.O. 1.2
CONSIGLIO NAZIONALE DELLE RICERCHE
ISTITUTO PER LA GENESI E L'ECOLOGIA DEL SUOLO - FIRENZE

# NORTHERN ASSO RIVER BASIN (SIENA - ITALY). ANALYSIS OF LINEARS - LANDFORMS - GEOLOGY RELATIONSHIPS AND ANTHROPIC EFFECTS IN THE PERIOD 1976 - 1994

E. BUSONI & A. COLICA

Scale 1: 25 000

## LINEARS, LANDFORMS AND ANTHROPIC EFFECTS IN 1976




## LINEARS, LANDFORMS, GEOLOGY AND ANTHROPIC EFFECTS IN 1994




## GEOLOGY

Geologic map according to the Servizio Geologico d'Italia, Foglio 121- Montepulciano, scale 1:100.000, map (after Jacobacci & Malatesta, 1968; Stratigraphy: after Lazzarotto (1963); Jacobacci et all (1963); Bicocchi et all (1987); Berti et all. (1992).

- ST — Tuscan Series (*upper Trias – Oligocene*).
- F — Flysch facies (*Oligocene-Miocene*)
- ps — Marine clays and sandy clays with conglomerate lenses. (*Early - Middle Pliocene*).
- pc — Polygenetic conglomerates, with several relationships with the other Pliocene deposits.
- tr — Travertines (*Pleistocene – Holocene*).
- f — Stream terrace (*Holocene*).
- e — Eluvial and colluvial deposits (*Holocene*).
- al — Alluvial deposits (*Holocene*).
- dt — Slope-debris (*Holocene*).

- dip
- normal fault
- normal buried fault
- anticline axis
- Linear: mainly recognized by aero-photo interpretation (photo scale: about 1:13000)

## LANDFORMS

(Landforms classification according with main morphogenetic processes)

### Erosional processes:

- **Biancana (Bi)**: a dome-shaped landform, usually present in clusters on hill slopes not steeper than 25%. Generally lower than 20 m, with a sub-horizontal pediment (or erosion-deposition glacis - *Castiglioni, 1979; Terni et all, 1994*). This landform usually develops in clay sediments (Ligurian). 1978- *Colica & Guaspari, 1990; Guaspari, 1993*. Biancane of similar form and size can also be found as clay knolls in pa, the dome has a vegetated (northern slope) or fully vegetated. In these two latter cases soils (from Entic to Inceptisols) are present. Rhomboidal nets of joints are related with the development of biancane; net joints directions may show linears directions (*Colica & Guaspari, 1990; Brondi et all, 1992; Bervegno et all, 1993*).

  position on the slope:
  top a
  mid m
  foot b

  tipology
  dome (after Calzolari, 1998) d
  vegetated V
  bare n
  vegetated only on a slope 1/2 V
  orthogonal direction to the main jua
  if the landform is circular the orthogonal direction to the main axis cannot be determined

- **Runoff surface:** surface created by the part of the rainfall that flows off, in non-concentrated form or in temporary, very shallow, channels (*Bergsma et all, 1996*).
- **Gully:** an incised channel or miniature valley cut by concentrated but intermittent runoff. It is deep enough to interfere with, and not to be obliterated by, normal tillage operations (*Bergsma et all, 1996*).
- **Shallow gully:** gully, not deeper and/or wider than 0.50 m, usually developed on sub-horizontal surfaces at the head of a calanco (*Busoni et all, 1998*).
- **Fluvial terrace scarp:** the landform derives from the cut of old fluvial surfaces, graded to the present base level (*Castiglioni, 1979; Soil Survey Staff, NRCS, 2001*).

### Erosional and gravitational processes (values in meters were retrieved on the 1:10000 base map)

- **Calanco (C):** small drainage basin, formed by small tallwegs separated by narrow interfluves (more or less sharp ridges with slope angles depending on the physical and mechanical characteristics of the bedrock (*Rodolfi & Frascati, 1979; Mazzanti & Rodolfi, 1988*). The landform develops on steep slopes (> 25%), mainly on clayey sediments, or clayey sediments stratified with textures coarser than clay (*Colica, 1992, 2000; Busoni & Colica, 2001*). It is generally more than 20 m high.

  If the calanco headcut is almost linear and the calanco is asymmetrical with respect to the main hydrographic axis:
  length (meters)
  width (meters)
  If the calanco headcut is arcuate and the calanco is symmetrical with respect to the main hydrographic axis:
  length (meters)
  length is the distance from head to the chord linking the its lowest points (meters)

- **Calanco potential area (CPA** *Busoni et all, 1998*): morphologically complex hill slope, generally steeper than 25%. It may represent either a transitional stage from a grazed or a cultivated area to a calanco, either a transition from a calanco area to a permanent, vegetated one. In the first case, features of erosional hot-spot processes is a calanco are evident; in the second case the calanco features are still recognizable, notwithstanding the presence of vegetation.

### Gravitational processes (values in meters were retrieved on the 1:10000 base map)

- **Shallow seated gravitational deformation (SSGD**, *Busoni et all, 1998*): elliptical, convex landform developed in pa, intergraded or not by sand layers, on hill slopes less steep than 20%; its main axis is shorter than 60 m but longer than that of biancane. Generally it is in clusters. It does not seem to be evidently related to a crown. Rhomboidal joint nets are related with the development of SSGD (*Colica & Guaspari, 1990*). Net joints directions may show linears directions.

  position on the slope
  top a
  mid m
  foot b
  extention of the main axis (meters)
  orthogonal direction to the main axis

- **Rock fall (For):** it develops in ps, pc-pc, ST and in sand lenses present in pa. It is characterized by a vertical or sub-vertical scarp. The scarp may coincide with joints.
- **Earthflow (Fco):** landform characterized by downslope translation of soil and weathered rock over a discrete basal shear surface within well defined lateral boundaries (*Soil Survey Staff, NRCS, 2001*). The crown may coincide with joints. This convex landform usually has dimensions larger than those of the SSGD. It develops in clay sediments, with or without sand interlayers.
- **Balza (Bz):** nearly vertical cliff, several meters high, mainly developed in arenites and rudites.
  linear l
  arcuate ar
  chord length (meters).

### Anthropic processes

- **Levelled biancana (Bl**, *Busoni et all, 1998*): the landform has been levelled because of the agricultural use of the slope. Its elliptical, or circular, base section is still visible.
- **Levelled shallow seated gravitational deformation (SSGDl**, *Busoni et all, 1998*): the landform has been levelled because of agricultural use. Its elliptical base section is still visible.

## NOTE

Landscapes change with time as a consequence of environmental state factor, anthropic included, changes. In fragile environments the changing of anthropic pressure, at present coincident with climate change, on landscape can be terrific and give origin to strong environmental degradation. Pressures on meta-stable state factors can trigger degradational processes by exceeding the threshold of resistance. Once the threshold is exceeded, a new state is reached. It is clear the importance of how and how much an environment changed in a specific time period is of paramount importance to build up scenarios and trends for the future. Most important is to control how anthropic pressures can lead to an increase in land degradation as a consequence of the influences of over-stressed landforms on hillslopes and watersheds hydraulics. Linear tectonic features are to be taken into account to define their influence on the evolution of a landscape and its landforms. The present maps highlight the changes occurred in the period 1976-1994 in a study area of central Italy. Linears and landforms (see Legend) were investigated. The analysis was carried on by aero-photo interpretation and field controls, taking into account linears, landforms, geology. Geology was used as first discriminant in the analysis of linears and landforms features. Data were reported on a 1:10000 scale base map and analysed on 1 km$^2$ base. Several linears were found to define directional trends showing the main regional faults directions: N(0°; N162°-N140°; N90°; N-S); these trends are called "main trends".

Tab. 1: Number of "main trends" linears relationship with geology (symbols as in legend).

| Geology | Main Trends | | | | |
|---|---|---|---|---|---|
| | (0°) | (162°) | (140°) | (90°) | (0°) |
| ps | 15 | 7 | 26 | 121 | 9 |
| pa | 5 | 7 | 9 | 9 | |
| ps pc | 2 | | | 1 | |
| ps ST | 9 | | | 1 | 5 |
| pa ps | | 1 | | | 1 |
| ST | 5 | 4 | 3 | 21 | |
| ST ps | | | | 1 | |
| ST ps pc | | | | | 5 |
| ST | | | | 1 | 1 |
| total | 36 | 25 | 32 | 165 | 9 |

Some other linears show directions different from the "main trends" ones: these linears are called "local linears". Different "linears landscapes" were found as a function of "local linears" density (= nr of linears/km$^2$), linears directions (°) and geology.




Fig. 1: Frequencies of "local linears" (number of linears/ km$^2$) and "linears landscapes". Nine linears landscape (fig. 1) were recognized among coincident with the 1:100000 Tuscany regional technical map sections covering the study area (NW: Acciano; N: S. Gimignano; NE: Sac. 258/130; W: Chiusure; Central: S. Giovanni d'Asso; E: Trequanda; SW: Montegiffoli; S: Castelmuzio; SE: Montefollonico).

Tab. 2: Local linears directional analysis in the nine "linears landscapes". Rayleigh's uniformity test calculates the probability p of the null hypothesis that the data are distributed in a uniform manner. A probability less than your chosen significance level (0.05) indicates that the data are not distributed uniformly and that they show evidence of a preferred direction.

| "Linears landscapes" | NW | N | NE | W | C | E | SW | S | SE |
|---|---|---|---|---|---|---|---|---|---|
| Observations | 110 | 220 | 74 | 191 | 378 | 133 | 33 | 214 | 31 |
| Mean vector (μ) | 85.42° | 59.38° | 141.79° | 49.79° | 63.34° | 48.80° | 83.72° | 69.47° | 55.70° |
| Length of mean vector (r) | 0.22 | 0.41 | 0.09 | 0.34 | 0.36 | 0.1 | 0.23 | 0.28 | 0.24 |
| Concentration | 0.45 | 0.9 | 0.17 | 0.72 | 0.77 | 0.19 | 0.47 | 0.65 | 0.5 |
| Circular variance | 0.78 | 0.59 | 0.92 | 0.66 | 0.64 | 0.9 | 0.77 | 0.74 | 0.76 |
| Circular standard deviation | 99.64° | 78.47° | 127.86° | 84.58° | 82.08° | 124.13° | 99.51° | 95.65° | 96.08° |
| Standard error of mean | 17.41° | 8.36° | 55.71° | 10.16° | 5.62° | 36.63° | 30.51° | 10.35° | 29.62° |
| Rayleigh test of uniformity (p) | 0.01 | 0 | 0.6 | 0 | 0 | 0.3 | 0.18 | 0 | 0.19 |

Tab. 3: Watson's test probability p matrix on "linears landscapes" directions. The p value printed for each test is the probability associated with the null hypothesis that the two mean angles are equal. If this probability is less than the chosen significance level (0.05), then the null hypothesis can be rejected in favour of the hypothesis that the two means are different. (red italics bold values)

| | NW | N | NE | W | C | E | S | SW | SE |
|---|---|---|---|---|---|---|---|---|---|
| NW | | 0.3 | 0 | 0 | 0.7 | 0.1 | 0.5 | 0.15 | 0.4 |
| N | | | 0 | 0.5 | 0.3 | 0.2 | 0 | 0.28 | 0.7 |
| NE | | | | 0 | 0 | 0 | 0 | 0 | 0 |
| W | | | | | 0 | 0.9 | 0 | 0 | 0.6 |
| C | | | | | | 0.3 | 0.2 | 0.04 | 0.4 |
| E | | | | | | | 0 | 0 | 0.6 |
| S | | | | | | | | 0.17 | 0.2 |
| SW | | | | | | | | | 0.1 |
| SE | | | | | | | | | |

Tab. 4: number of "local linears" and their percentage as function of geology, in each landscape (symbols as in legend).

| | | | |
|---|---|---|---|
| NW | linears: 110 | N | linears: 220 |
| pa: 97.3; ps: 0.9; tr: 1.8 | | pa: 63.6; pc: 34.1; ST: 0.9; tr: 0.5; al: 0.9 | |
| W | linears 131 | Central | linears: 378 |
| pa: 99.2; ps: 0.0; pc-pc: 0.8 | | pa: 80.7; pc: 0.3; ST: 19.0 | |
| SW | linears: 33 | S | linears: 214 |
| pa: 87.9; ps: 0.0; al: 12.1 | | pa: 92.0; ps: 5.0; al: 3.0 | |
| NE | linears: 74 | | |
| pa: 1.4; ps: 98.6 | | | |
| E | linears: 133 | | |
| pa: 0.0; ps: 0.0; ST: 100.0 | | | |
| SE | linears: 31 | | |
| pa: 0.0; ps: 0.0; pc-pc: 100.0 | | | |

Tab. 5: landforms relationships with geology (symbols as in legend).

| Landforms | Geology | | | |
|---|---|---|---|---|
| | pa | ps | ps - pc | ST |
| C | + | | | |
| CPA | + | | | |
| Bi - Bl | + | | | |
| SSGD - SSGDl | + | | | |
| Fco | + | | | |
| For | | + | + | |
| Bz | | | | + |

Tab. 6: anthropic effect. Variations in landforms presence in the period 1976 – 1994, and their relationships with linears. It is to be considered that most of the differences are effects of anthropic pressures on the environment. Decrease in levelled landforms must be seen as obliteration of their base sections by land use; these cannot be recognized neither by photo-interpretation, nor by field survey; the same should be held for their relationships with linears (symbols as in legend).

| | 1976 | | | 1994 | | | Differences between totals | | |
|---|---|---|---|---|---|---|---|---|---|
| | Total nr of landforms | Related to linears | Not related to linears | Total nr of landforms | Related to linears | Not related to linears | Abs. value | % | |
| | | % | % | | % | % | | | |
| C | 309 | 76.1 | 23.9 | 201 | 76.1 | 23.9 | -108 | -34.9 | |
| CPA | 54 | 40 | 74.1 | 14 | 28.6 | 57 | 36 | 63.2 | 21 | 36.8 | +3 | +5.6 |
| Bi | 54 | 38 | 70.4 | 16 | 29.6 | 40 | 24 | 60.0 | 16 | 40.0 | -14 | -25.9 |
| Bl | 57 | 26 | 45.6 | 31 | 54.4 | 18 | 18 | 100 | | | -39 | -68.4 |
| SSGD | 30 | 30 | 100 | | | 15 | 7 | 46.6 | 8 | 53.4 | -15 | -50 |
| SSGDl | 156 | 66 | 54.5 | 71 | 45.5 | 71 | 47 | 66.2 | 24 | 33.8 | -85 | -54.5 |
| Fco | 77 | 19 | 70.4 | 8 | 29.6 | 19 | 19 | 100 | | | -8 | -29.6 |
| For | 87 | 55 | 63.2 | 32 | 36.8 | 65 | 52 | 80.0 | 13 | 20.0 | -22 | -25.3 |
| Bz | 6 | 4 | 66.6 | 2 | 33.4 | 1 | 0 | 0 | 1 | 100 | -5 | -83.3 |

### Typology of the relationships between linears and landform.

Linears can relate to landforms in several ways. The following relationships were defined and shown in Tables 5 and 6:

l - linears limit the landform. (mainly C and CPA);
ac - linears coincide with the main hydrographic axis representing the base level for tributaries which head in the landform (mainly C and CPA);
l-head - linears coincide with the landform head (mainly C and CPA);
ad - linears coincide with main hydrographic axis; direct effect (mainly C and CPA);
t - linears cut the landform, not coinciding with hydrographic axis (mainly C and CPA);
d - linears participate to the net in which the landform develops (mainly Bi, Bl and SSGD, SSGDl);
p - linears are parallel to the crest of the landform (mainly For, usually in pa, pc, ST);
cl - linears are parallel to the landform;
al - linears are aligned with the landform (mainly For and Bz, usually in pa, pc, ST).

Tab. 7: typology percentage of the relationships between linears and landform in pa (symbols as in legend).

| Landform | | | | | | |
|---|---|---|---|---|---|---|
| | l | ac | l-head | ad | t | cl |
| C | 24.3 | 15.8 | 4.5 | 10.2 | 45.2 | - |
| CPA | 8.8 | - | 8.8 | 52.9 | 29.5 | - |
| Bi - Bl | - | - | - | - | - | 100 |
| SSGD - SSGDl | - | - | - | - | - | 100 |
| Fco | - | - | - | 100 | - | - |

Tab. 8: typology percentage of the relationships between linears and landform in pa, pc, st (symbols as in legend).

| Landform | | | | | | |
|---|---|---|---|---|---|---|
| | l | ac | t | d | other | al |
| For | 14.3 | 28.6 | 14.3 | 28.6 | 7.1 | 7.1 |
| Bz | | 50 | | 25 | | 25 |

Note to tables: data in the tables refer to real values of the different variables analized, as counted on the working map at the 1:10000 scale, on which photo-interpretation data were reported. The presented maps, at the 1:25000 scale, are the original to the 1:25000 scale; some of the original features were deleted to keep clearness in the readability of the maps. This is the reason why sometimes the area of study is comparison between landforms in the period 1976-1994, so that data in the tables slightly differ from map data.

### References

BENVEGNU F., BRONDI A., COLICA A., CONTI P., GUASPARI G., POLIZZANO C., SANNA G. & TASSONI E. (1993) - *Studies of regolith factors in clay shale in real situations. Study of fractures in clays of the neogenic basin of Siena*. Nuclear Science and Technology Series, European Communities Commission, Luxembourg.Office for Official Publications of European Communities, pp. 53.

BERGSMA E., CHARMAN P., GIBBONS F., HURNI H., MOLDENHAUER W.C., RINNCHPOROS S. (1996) - *Terminology for soil erosion and conservation*. ISSS, Wageningen, pp. 313.

BERTI A., CALAMERI G., CALITERNA L., FORLANI CONTI M., FARSI F., GUIDI F., LAZZAROTTO A., MANTOANI E., MICHELUCCI M., NEZZI GRIFI L. (1992) - *Il Duomo di Pienza - 1549 - 1994. Studi e Ricerche*. Ministero per i Beni Culturali ed Ambientali, Soprintendenza per i Beni Ambientali ed Architettonici di Siena. Ed. Centro, Firenze, pp. 167.

BOCCALETTI M., DECANDIA F.A., GASPERI G., GELMINI R., LAZZAROTTO A., ZANZUCCHI G. (1987) - *Note illustrative, Carta strutturale dell'Appennino settentrionale*. CNR - PF. Geodinamica, Sottoprog. 3 Modelli strutturali, Gruppo Appennino settentrionale, Tip. Senese, pp. 203.

BRONZI A. & COLICA A., CONTI P., GUASPARI G, POLIZZANO C., SABATINI G. (1992) - *Joints in clay of neogenic basin of Siena*. Mineralogica et Petrographica Acta, XXXV, Bologna, 51-65.

BUSONI E., COLICA A. (2001) - *La rivalorizzazione dell'habitat integrata del paesaggio: loro relazioni con forme di degradazione del suolo. Atti dal Convegno Nazionale S.I.S.S., Venezia 10-16 giugno 2000*, Boll. Soc. It. Scienza del Suolo, Vol. 50: 251-260.

BUSONI E., COLICA A., SALVADOR SANCHES P. (1998) - *Integrated analysis of the Arno river basin (Siena Italy)*. C.N.R. Institute for Soil Genesis and Ecology, Florence, 1998 (Map, scale 1:30000).

CALZOLARI C., UNGARO T. (1998) - *Geomorphic features of a badland (biancana) area (Central Italy): characterization, distribution and quantitative spatial analysis*. CATENA, 31, 237-256.

CASTIGLIONI G.B. (1979) - *Geomorfologia*. UTET, Torino, pp. 436.

COLICA A. (1992) - *Processi erosivi in calanchi e biancane e loro implicazioni per le opere di conservazione nell'alta Val d'Orcia (SI)*. Tesi di dottorato di ricerca, Università degli Studi di Firenze, pp. 165.

COLICA A. (2000) – *Analysis of the genesis and evolution of calanco morphology in an experimental area of Piazza (Radicofani, Siena, Italy)*. Geogr. Fis. e Dinam. Quat. 23.

COLICA A. & GUASPARI G. (1990) - *Sistemi di fratturazione nelle argille plioceniche del territorio senese. Implicazioni geomorfologiche*. Atti Accad. Fisiocritici Siena, Serie IV, Torno IX, 29-36.

GUASPARI G. (1978) - *Calanchi e biancane nel territorio senese: studio geomorfologico*. L'Universo, 48, 97-140.

GUASPARI G. (1993) - *I rivestimenti geomorfologici dei terreni argillosi pliocenici*. In: La storia naturale della Toscana meridionale, M.P.S., 89-106.

JACOBACCI A., MALATESTA A. (1968) - *Carta Geologica d'Italia. Montepulciano, Foglio 121*. Scala 1:100.000. G.N.M.

JACOBACCI A., MALATESTA A., MARTELLI G. (1963) *Note illustrative della Carta Geologica, foglio 121 – Montepulciano*. Servizio Geologico d'Italia, pp. 73.

LOSACCO U. (1963) - *Osservazioni geologiche e geomorfologiche nei dintorni di Pienza e nella media Val d'Orcia*. L'Universo, 48, 4, 659-688.

MAZZANTI R. & RODOLFI G. (1988) - *Evoluzione del rilievo nei sedimenti argillosi a stabilità di cliff retrogressiva e quaternari ballani*. In: Canuti P. & Pranzini E. (Editors): la gestione delle aree fragili. Ed.delle Autonomie, 13 - 60.

RODOLFI G. & FRANZOSI F. (1979) - *Carta geologica di base per la programmazione degli interventi in aree marginali*. Annali Ist. Spec. Studio e Difesa del Suolo, X, Firenze, 38 - 80.

SOIL SURVEY STAFF, NRCS (2001) - *National Soil Survey Handbook*. Title 430 - VI, Part 629.

TORRI D., COLICA A., ROCKWELL D. (1994) - *Preliminary study of erosion mechanics in a biancane badland (Tuscany - Italy)*. Catena, Catena Verlag, Braunschweig (Germania), 1994, 281-294.

*Acknowledgements*

The research was supported by CNR – GNDCI grants.
The map is published with funds from the CNR – GNDCI Presidency.

*Base topografica:*
Regione Toscana - Carta Topografica 1: 25 000 da fpt dell'istituto Geografico Militare (sul. n. 2401 in data 18-03-1996)
Regione Toscana - Ortofotocarta 1: 10 000 Rilievo: aereo 1977- Ed. alle stuttazione: SMA – N. 56 in data 21-02-1997

*Prepared and printed by:* S.EL.C.A. via Reginaldo Giuliani, 153 - Firenze - 2002